PREMIER ORAL FACIAL & IMPLANT SURGERY FIRAS ALI, DMD

PRE-OPERATIVE INSTRUCTION FOR SURGERY

PRE-OPERATIVE INSTRUCTION FOR SURGERY

- 1. Any patient under 18 years of age must be accompanied by parent or guardian at the time of surgery.
- 2. Please wear loose fitting clothing with sleeves that can be rolled up past the elbow and low heel shoes.
- 3. Take all your regular medication with a sip of water unless you are instructed to do otherwise by the doctor in this office or your primary medical doctor.
- 4. IF YOU ARE SCHEDULED TO HAVE <u>INTRAVENOUS SEDATYION</u>: you may not have anything to EAT or DRINK FOR 8 HOURS before your appointment. For morning surgery no food or liquids after midnight the night before surgery. Clear liquids may be be taken up to 2 hours before surgery.
- 5. If you are scheduled to have Nitrous Oxide Analgesia (N2O), you may not have anything to eat or drink including water for 2 hours before appointment.
- 6. NO smoking for at least 12 hours before surgery. Ideally cut down or stop smoking as soon as possible before the day of surgery.
- 7. A responsible Adult MUST accompany the patient to the office, remain in the office durig the procedure and drive patient home.
- 8. Plan to rest the reminder of the surgery day. The Patient should not DRIVE a vehicle, operate any machinery or make important decisions for 24 hours following Intravenous Anesthesia.
- 9. DO NOT wear lipstick excessive makeup or nail polish on the day of surgery.
- 10. Contact lenses, jewelry and dentures MUST be removed before surgery.
- 11. The use of alcohol or recreational drugs can have an adverse effect on the anesthesia medication that we use. Please discontinue the use of such drugs for at least 72 hours before procedure.
- 12. If you have an illness such as Cold, Sore Throat or Stomach upset please notify the office.
- 13. If for any reason you are unable to keep your surgery appointment, we request a 48 hours notice of cancellation.
- 14. Women, please note: some antibiotics may interfere with the effectiveness of your birth control pills, please check with your pharmacist.